

## Estimated 1-5 Year Cancer Survivors 2011

STATE	2,011 Estimate Cases	1 Year Survival	2,010 Estimate Cases	2 Year Survival	2,009 Estimate Cases	3 Year Survival	2,008 Estimate Cases	4 Year Survival	2,007 Estimate Cases	5 Year Survival	2,006 Estimate Cases	Estimated 1-5 Year Survivors
AL	25,530	76.8%	23,640	68.4%	24,090	63.3%	22,340	59.5%	20,590	56.3%	24,390	74,760
AK	3,090	76.8%	2,860	68.4%	2,530	63.3%	2,650	59.5%	2,500	56.3%	2,010	8,220
AZ	31,550	76.8%	29,780	68.4%	27,600	63.3%	25,540	59.5%	26,270	56.3%	25,450	87,880
AR	16,070	76.8%	15,320	68.4%	14,800	63.3%	14,840	59.5%	14,130	56.3%	15,220	48,260
CA	163,480	76.8%	157,320	68.4%	152,170	63.3%	156,530	59.5%	151,250	56.3%	138,680	492,060
CO	22,390	76.8%	21,340	68.4%	20,340	63.3%	18,900	59.5%	19,190	56.3%	16,850	63,170
CT	21,440	76.8%	20,750	68.4%	20,650	63.3%	19,190	59.5%	19,780	56.3%	17,320	63,730
DE	5,130	76.8%	4,890	68.4%	4,690	63.3%	4,590	59.5%	4,530	56.3%	4,190	14,920
DC	2,830	76.8%	2,760	68.4%	2,600	63.3%	2,560	59.5%	2,540	56.3%	2,680	8,540
FL	113,400	76.8%	107,000	68.4%	102,210	63.3%	101,920	59.5%	106,560	56.3%	98,960	335,720
GA	44,580	76.8%	40,480	68.4%	39,080	63.3%	36,980	59.5%	35,440	56.3%	36,650	122,950
HI	6,710	76.8%	6,670	68.4%	6,400	63.3%	6,310	59.5%	6,020	56.3%	5,600	20,230
ID	7,520	76.8%	7,220	68.4%	6,800	63.3%	6,430	59.5%	6,140	56.3%	5,900	21,240
IL	65,610	76.8%	63,890	68.4%	60,960	63.3%	59,130	59.5%	62,010	56.3%	60,220	198,990
IN	34,050	76.8%	33,020	68.4%	31,320	63.3%	29,550	59.5%	30,040	56.3%	32,710	101,780
IA	17,500	76.8%	17,260	68.4%	16,740	63.3%	16,150	59.5%	16,540	56.3%	16,310	53,950
KS	14,070	76.8%	13,550	68.4%	13,080	63.3%	12,520	59.5%	12,760	56.3%	13,210	42,310
KY	25,010	76.8%	24,240	68.4%	24,060	63.3%	23,270	59.5%	22,850	56.3%	23,690	76,740
LA	22,780	76.8%	20,950	68.4%	22,170	63.3%	23,360	59.5%	22,540	56.3%	23,890	72,900
ME	8,820	76.8%	8,650	68.4%	9,000	63.3%	8,140	59.5%	8,340	56.3%	7,910	27,370
MD	28,890	76.8%	27,700	68.4%	26,650	63.3%	27,380	59.5%	26,390	56.3%	25,870	87,100
MA	37,470	76.8%	36,040	68.4%	36,080	63.3%	34,470	59.5%	34,920	56.3%	33,310	113,710
MI	57,010	76.8%	55,660	68.4%	53,550	63.3%	51,150	59.5%	54,410	56.3%	48,250	171,290
MN	27,600	76.8%	25,080	68.4%	23,670	63.3%	23,160	59.5%	25,420	56.3%	23,520	78,480
MS	14,990	76.8%	14,330	68.4%	14,150	63.3%	13,400	59.5%	12,470	56.3%	15,120	45,100
MO	32,740	76.8%	31,160	68.4%	30,090	63.3%	29,390	59.5%	29,930	56.3%	31,050	98,410
MT	5,690	76.8%	5,570	68.4%	5,340	63.3%	5,090	59.5%	4,920	56.3%	4,780	16,770
NE	9,430	76.8%	9,230	68.4%	8,810	63.3%	8,710	59.5%	8,720	56.3%	8,450	28,570
NV	12,800	76.8%	12,230	68.4%	12,020	63.3%	11,370	59.5%	11,030	56.3%	11,600	37,910
NH	8,210	76.8%	7,810	68.4%	7,630	63.3%	7,030	59.5%	7,140	56.3%	6,470	23,560
NJ	49,080	76.8%	48,100	68.4%	47,920	63.3%	45,900	59.5%	49,370	56.3%	43,910	152,870

STATE	2,011 Estimate Cases	1 Year Survival	2,010 Estimate Cases	2 Year Survival	2,009 Estimate Cases	3 Year Survival	2,008 Estimate Cases	4 Year Survival	2,007 Estimate Cases	5 Year Survival	2,006 Estimate Cases	Estimated 1-5 Year Survivors
NM	9,630	76.8%	9,210	68.4%	8,830	63.3%	8,260	59.5%	8,030	56.3%	8,150	27,710
NY	107,260	76.8%	103,340	68.4%	101,550	63.3%	97,130	59.5%	100,960	56.3%	88,230	320,050
NC	48,870	76.8%	45,120	68.4%	42,270	63.3%	40,420	59.5%	38,210	56.3%	40,890	134,910
ND	3,560	76.8%	3,300	68.4%	3,200	63.3%	3,090	59.5%	3,340	56.3%	3,170	10,450
OH	65,060	76.8%	64,450	68.4%	62,420	63.3%	56,840	59.5%	59,220	56.3%	61,310	197,930
OK	18,980	76.8%	18,670	68.4%	18,110	63.3%	17,860	59.5%	17,170	56.3%	18,640	58,740
OR	21,180	76.8%	20,750	68.4%	19,210	63.3%	19,230	59.5%	18,630	56.3%	18,290	62,630
PA	78,030	76.8%	75,260	68.4%	74,170	63.3%	70,110	59.5%	75,130	56.3%	73,630	239,070
RI	6,090	76.8%	5,970	68.4%	6,250	63.3%	6,120	59.5%	6,360	56.3%	5,900	19,840
SC	25,510	76.8%	23,240	68.4%	22,100	63.3%	20,740	59.5%	21,370	56.3%	22,530	71,490
SD	4,430	76.8%	4,220	68.4%	4,120	63.3%	4,080	59.5%	3,990	56.3%	4,060	13,300
TN	34,750	76.8%	33,070	68.4%	32,570	63.3%	29,390	59.5%	28,440	56.3%	32,140	101,300
TX	105,000	76.8%	101,120	68.4%	98,200	63.3%	96,320	59.5%	91,020	56.3%	85,570	308,130
UT	10,530	76.8%	9,970	68.4%	8,880	63.3%	7,760	59.5%	7,660	56.3%	6,620	26,930
VT	3,950	76.8%	3,720	68.4%	3,550	63.3%	3,530	59.5%	3,500	56.3%	2,970	11,270
VA	38,720	76.8%	36,410	68.4%	34,150	63.3%	35,590	59.5%	35,090	56.3%	34,990	114,430
WA	35,360	76.8%	34,500	68.4%	32,290	63.3%	32,380	59.5%	31,080	56.3%	28,350	103,530
WV	11,080	76.8%	10,610	68.4%	10,230	63.3%	10,250	59.5%	10,490	56.3%	11,280	34,230
WI	30,530	76.8%	29,610	68.4%	27,560	63.3%	27,590	59.5%	28,130	56.3%	26,390	90,650
WY	2,680	76.8%	2,540	68.4%	2,500	63.3%	2,570	59.5%	2,340	56.3%	2,500	8,090

Survival is assumed to be similar across states and is based on observed survival among cases diagnosed in SEER 17 areas during 1999-2007, followed through December 2008. The Estimated 1-5 Year Survivors do not include 2011 cases because survival is defined here as those surviving at least 1 year; the number of 1-5 year survivors is computed by the summation of 5 products: 2006 estimated cases multiplied by the 5-year observed survival rate, 2007 estimated cases multiplied by 4 year survival rate, 2008 estimated cases multiplied by 3 year survival rate, 2009 estimated cases multiplied by 2 year survival rate, and 2010 estimated cases multiplied by the 1 year survival rate.